

Pregnancy outcomes in women after an arterial switch operation for transposition of the great arteries

Department of Obstetrics and Gynecology¹⁾

Department of Pediatrics²⁾

National Cerebral and Cardiovascular Center, Osaka, Japan

Chinami Horiuchi¹⁾, Chizuko Kamiya¹⁾, Atsushi Daimon¹⁾, Masami Sawada¹⁾,

Tetsuya Ide¹⁾, Takekazu Miyoshi,¹⁾Rina Funada¹⁾, Mitsuhiro Tsuritani¹⁾,

Naoko Iwanaga¹⁾, Reiko Neki¹⁾, Jun Yoshimatsu¹⁾, Hideo Ohuchi²⁾, Isao Shiraishi²⁾

Introduction

- The arterial switch operation (ASO) has replaced the atrial switch operation and has become the standard operation for improving the survival and quality of life of patients with TGA.
- Many women with TGA, wishing to have children, are now expected to reach childbearing age.
- The purpose of this study was to determine the risk of a pregnancy with TGA repaired by ASO.



Methods

Study design : retrospective cohort study

Population : 11 pregnancies in 8 women after ASO
between 2006-2015 at our hospital

Reviews of records

Clinical status, echocardiography, and plasma BNP level

Main outcomes

Adverse maternal cardiovascular events

- arrhythmia : need medical therapy
- heart failure : lower oxygen saturation with pulmonary edema
- deteriorated NYHA function class
- coronary event: chest pain and/or ST change in electrocardiography

Obstetrical events

Neonatal events

Results

Baseline clinical characteristics

Type	Prior palliation	Age at ASO	Re-intervention Since ASO	NYHA	Cardiac finding	Arrhythmia	Coronary artery disease	Medication	
A	2	—	11M	Right -sided	I	PR mod	-	-	-
B	1	BAS	3M	—	I	PS mod	-	-	-
C	1	BAS, PAB, ltBTS PDA ligation	6M	—	I	-	-	-	-
D	2	BAS PAB, lt BTS	11M	—	I	Aortic root dilation (54mm) AR mod, PR mod	-	-	Carvedilol
E	2	BAS	12M	Left -sided	I	MR mod, AR mod residual VSD	-	-	Diuretics
F	2	BAS	5M	Right -sided	I	-	-	-	-
G	1	BAS	6D	Right -sided	I	-	-	-	-
H	1	BAS	15D	Right -sided	I	-	-	-	-

A, B, C, E complicated with cardiovascular events in pregnancy and/or postpartum

Obstetric outcomes / events


Mean maternal age (years)	25.3 ± 5.3
Miscarriage	0
Parity status 1 / 2 / 3	5 / 2 / 1
Twin pregnancy	1
Obstetric events	
Pregnancy-induced hypertension	1 (9%)
Threatened premature delivery	1 (9%)
Fetal growth restriction	2 (18%)
Postpartum hemorrhage	1 (9%)
Mean gestational week of delivery (weeks)	37.6 ± 1.8
Mode of delivery	
Vaginal	6 (55%)
Cesarean section	5 (46%)
Indication: Cardiac / Obstetric	2 (40%) / 3 (60%)

Neonatal Outcomes in 12 babies

Neonatal birth weight (g) 2598 ± 616

Neonatal events

Preterm birth (<37weeks)	3 (25%)
Very preterm birth (<34weeks)	0
Small - for-gestational age	4 (36.4%)
Heavy -for-gestational age	1 (8.3%)
Transient tachypnea of neonate	2 (16.7%)
Congenital heart disease	0







Overview of pregnancies with cardiovascular events

	Gestational Weeks	Event	Treatment
A	24 postpartum	NSVT increased PVC	Start Carvedilol Carvedilol ↑
B	37	NSVT	rest (asthma +)
C	post	Pulmonary edema	Diuretic
	28 31	NSVT Atrial tachycardia	Start Carvedilol Carvedilol ↑
E	post	Pulmonary edema NSVT	Diuretics ↑ Carvedilol ↑ ACE-I

Case E

twin pregnancy

Serial BNP levels and the cardiovascular events



Risk factors for cardiovascular events

	Event (+) (n=4)	Event (-) (n=7)	P-value
TGA type 1	2	6	
type 2	2	1	ns
Re-operation post ASO	2	5	ns
Cardiac medications before pregnancy	1	1	ns
Twin pregnancy	1	0	ns
Tricuspid valve regurgitation $PG \geq 30$ mmHg during pregnancy	4	1	0.015
Aortic valve regurgitation (moderate/severe)	1	1	ns
Neo-aortic root dilatation (> 40mm)	2	1	ns
Pulmonary valve regurgitation(moderate/severe)	1	1	ns
Left ventricular end-diastolic diameter(≥ 50 mm) during pregnancy	4	2	ns
BNP ≥ 30 pg/ML in 1 st trimester	4	1	0.015

Serial TRPG and BNP levels in pregnancies with and without cardiovascular events



—◆— Event (+)
—■— Event (-)

TRPG

BNP

(mmHg)

(pg/ML)



※ P < 0.05 between red and blue

Long-term follow-up after pregnancy (3– 120 months)

	No. Pregnancy	NY HA	Coronary artery disease	PS		PR		AR		Aortic root dilatation	
				before	after	before	after	before	after	before	after
A	1	I	-	-	-	Mod.	Severe	-	-	-	-
B	1	I	-	Mod.	Mod.	-	-	-	-	-	-
	2	I	-	Mod.	Mod.	-	-	-	-	-	-
C	1	I	-	-	-	-	-	-	-	-	-
D	1	I	-	-	-	Mod.	Mod.	-	-	+	+
										54mm	54mm
E	1	I	-	-	-	-	-	Mod.	Mod.	-	-
G	1	I	-	-	-	-	-	-	-	-	-
H	1	I	-	-	-	-	-	-	-	-	-
	2	I	-	-	-	-	-	-	-	-	-
	3	I	-	-	-	-	-	-	-	-	-

- ; abscent or mild

Case F was excluded because the last examinaton was performed at 3 days postpartum

Discussion – Reviews of previous reports –

	Arterial switch operation		Atrial switch operation (Mustard/Senning)					
	Horiuchi 2016	Tobler 2010	Guédès 2004	Drenthen 2005	Metz 2011	Doninica 2012	Trigas 2014	Cataldo 2015
n (women/pregnacy)	8/11 (twin 1)	9/13	16 / 28	-/49 (twin 2)	10/14	19/43 (twin 1)	34/44	34/21
Cardiac events								
deteriorated NYHA	0		7	17	2/14		7	
Arrhythmia	3(27%)	1	2	11	1	2	4	
Heart failure	2 (18%)	1	2	2	1	2 Death 1	5 Arrest 1	
Increased of TR	0	-	7/20	-	+	-	-	-
systemic ventricle dysfunction	0	2 (valve thrombus 1)	4/21	33%	89%			
Obstetric events								
PE /PIH /HELLP	1		1	10	1	1	1	1
FGR	1							5
PE / DVT				2				
Neonatal events								
premature labor	3	0	-	7	7	8	11	10
SGA (< 10%)	4	1		11	-	-	5	10
IUFD	0	-		4	-	2	-	-
Long-term outcome								
valve lesions	PR↑ 1 (14%)	PS ↑ 1	TR ↑ 2		-			TR↑ (33%)
systemic ventricular dysfunction		1	4			2 Death 1		(67%)
NHHA(↓)	0	-	2	(8.2%)	-		5	(14%)

Conclusion

- ✓ Pregnancy after ASO was well-tolerated.
 - The most severe case was complicated with twin pregnancy; Such cases need special attentions.
- ✓ Neonatal outcomes were mostly good.
- ✓ Women with the RV hypertension or the higher plasma BNP levels in 1st trimester tended to have adverse cardiovascular events in the peripartum period.
- ✓ Pregnancy seems to have less adverse impact in women after ASO compared to women after atrial switch operation, who have the systemic right ventricle.